

Abstract

These notes are based on Ron Donagi's lectures for the course "Complex Algebraic Geometry" at UPenn along with Daniel Huybrechts's *Complex Geometry*. Any mistake in what follows is my own.

Contents

1 A cursory overview of algebraic geometry	2
1.1 Lectures 1-4	2
2 Complex analysis	4
2.1 Lecture 5	4
3 Line bundles	6
3.1 Lecture 6	6
3.2 Lecture 7	9
3.3 Lecture 8	11
3.4 Lecture 9	14
3.5 Lecture 10	16
4 Kähler manifolds	19
4.1 Lecture 11	19
4.2 Lecture 12	22
5 Lie algebras	24
5.1 Lecture 13	26

1 A cursory overview of algebraic geometry

1.1 Lectures 1-4

These lectures consisted of informal surveys of certain fundamental concepts of algebraic geometry. They were meant as previews of various topics that we will cover rigorously. The following three problems give a taste of the initial concepts presented.

Projective space.

Recall that projective space \mathbb{P}^n of dimension n is by definition the quotient of $\mathbb{A}^{n+1} \setminus \{0\}$ by the rescaling action of $\text{GL}(1)$. This is a compactification of n -dimensional affine space \mathbb{A}^n . Show that \mathbb{P}^n has an open covering by $n + 1$ many open subsets U_0, \dots, U_n each isomorphic to \mathbb{A}^n . Work out all of the transition functions.

For simplicity, assume that $\mathbb{A}^n = \mathbb{R}^n$. For each $i \in \{1, \dots, n + 1\}$, let $\widetilde{U}_i = \{\vec{x} \in \mathbb{R}^{n+1} : x_i \neq 0\}$. Let $\pi : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}\mathbb{P}^n$ denote the natural projection and let $U_i = \pi(\widetilde{U}_i)$. Since \widetilde{U}_i is saturated and open, we know that $\pi|_{\widetilde{U}_i}$ is a quotient map. Define $f_i : U_i \rightarrow \mathbb{R}^n$ by $[x_1, \dots, x_{n+1}] \mapsto \left(\frac{x_1}{x_i}, \dots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \dots, \frac{x_{n+1}}{x_i}\right)$, whose inverse is given by $(x_1, \dots, x_n) \mapsto [x_1, \dots, x_{i-1}, 1, x_{i+1}, \dots, x_n]$. Since $f_i \circ \pi$ is continuous, so is f_i . Hence we see that f_i is a homeomorphism.

Now, consider any transition map $f_i \circ f_j^{-1}$. Let $1 \leq i, j \leq n + 1$. If $i = j$, then the transition map is the identity, which is smooth. So assume, wlog, that $j < i$. Then

$$f_j \circ f_i^{-1}(x_1, \dots, x_n) = \left(\frac{x_1}{x_j}, \dots, \frac{x_{j-1}}{x_j}, \frac{x_{j+1}}{x_j}, \dots, \frac{x_{i-1}}{x_j}, \frac{1}{x_j}, \frac{x_i}{x_j}, \dots, \frac{x_n}{x_j}\right)$$

when $(x_1, \dots, x_n) \in f_i(\widetilde{U}_i \cap \widetilde{U}_j)$. This is clearly smooth. ■

Classification of conics.

Classify, up to linear automorphism, all conics (i.e., 1-dimensional quadratic hypersurfaces) in $\mathbb{R}\mathbb{P}^2$.

First of all, we may assume that any conic $C(x, y, z) = 0$ in $\mathbb{R}\mathbb{P}^2$ is a homogeneous polynomial of degree two. As a result, we can write $C(x, y, z) = v^t A v$ for some symmetric 3×3 matrix A over \mathbb{R} where

$$v := \begin{bmatrix} x \\ y \\ z \end{bmatrix}.$$

Since A is orthogonally diagonalizable, we have that $C(v) = (Tv)^t D(Tv)$ for some orthogonal matrix T and some diagonal matrix D . Thus, C is isomorphic to a curve in $\mathbb{R}\mathbb{P}^2$ defined by $0 = a_1x^2 + a_2y^2 + a_3z^3$. We can apply another invertible change of variables to ensure that $a_i \in \{-1, 0, 1\}$ for each $i = 1, 2, 3$.

Likewise, any quadratic hypersurface in $\mathbb{C}\mathbb{P}^n$ is isomorphic to a curve in $\mathbb{C}\mathbb{P}^n$ defined by $\sum_{i=1}^{n+1} a_i x_i^2 = 0$ where $a_i \in \{-1, 0, 1\}$ for each $i = 1, \dots, n + 1$.

We now see that there are exactly five distinct automorphism classes of conics in $\mathbb{R}\mathbb{P}^2$:

- (a) $[x^2 = 0]$ (a double line),

- (b) $[x^2 + y^2 = 0]$ (a single point),
- (c) $[x^2 - y^2 = (x - y)(x + y) = 0]$ (the union of two distinct lines that intersect at a single point),
- (d) $[x^2 + y^2 + z^2 = 0]$ (the empty set), and
- (e) $[x^2 + y^2 - z^2 = 0]$ (a curve C whose affine part is either an ellipse, a parabola, or a hyperbola, depending on whether C intersects the line at infinity at zero, one, or two points, respectively).

■

Plane cubics.

Recall that a plane curve C is *smooth* if it has no *singular* points, i.e., points at which all first-order partial derivatives of C vanish.

1. Consider a smooth plane cubic X with an inflection point, i.e., a non-singular point at which the tangent line to X has intersection multiplicity at least 3 with X . Show that X is isomorphic, up to linear automorphism of the ambient space, to a smooth plane cubic in *Weierstrass form* $y^2 = x^3 + ax + b$.
2. Find an expression of the form $J := \frac{a^m}{b^n}$ for some integers m and n that is invariant under linear changes of coordinates sending our equation of X to another Weierstrass form.
3. A slight problem with J is that it goes to infinity when $b = 0$ even though the plane cubic X remains smooth. Can you suggest another function $j(a, b)$ which has the same invariance property as J but remains finite (i.e. is holomorphic) whenever X is smooth? Relate j to J .

1. Any linear automorphism of \mathbb{C}^3 induces a projective transformation of \mathbb{CP}^2 (i.e., an element of $\text{PGL}(3, \mathbb{C})$), and the group of all linear automorphisms of \mathbb{C}^3 acts transitively on the set of all 2-dimensional subspaces of \mathbb{C}^3 . It follows that for any two lines $L_1, L_2 \subset \mathbb{CP}^2$, there is some projective transformation $\psi : \mathbb{CP}^2 \rightarrow \mathbb{CP}^2$ such that $\psi(L_1) = L_2$. In particular, we can assume that the tangent line L to our given inflection point P is precisely the line at infinity $\{z = 0\}$ and that any other line touching P is precisely the line $\{x = 0\}$. This means that P corresponds to the point $[0, 1, 0]$ on our new, isomorphic copy X' of X . Note that X' is smooth since any projective transformation preserves smoothness by the familiar multivariable chain rule.

Let X' be the zero locus of the homogeneous polynomial $f(x, y, z)$ of degree three. This has the form

$$Ax^3 + Bxy^2 + Cyx^2 + Dy^3 + zg(x, yz)$$

where $g(x, y, z)$ denotes a homogenous polynomial of degree two. We have that $d = 0$ because $[0, 1, 0]$ lies on X' . Further, we have that $b = c = 0$ because $\{z = 0\}$ is tangent to an inflection point. Therefore, the affine part of X' has the form

$$f(x, y, 1) = Ax^3 + Bx^2 + Cxy + Dy^2 + Ex + Fy + G$$

where both A and D are nonzero. By scaling this appropriately, we may write

$$f(x, y, 1) = x^3 + Bx^2 + Cxy - y^2 + Ex + Fy + G$$

for some $B, C, E, F, G \in \mathbb{C}$.

It remains to show that there exists an invertible affine map (or change of variables) $\mathbb{C}^2 \rightarrow \mathbb{C}^2$ that sends the zero locus of $f(x, y, 1)$ to the zero locus of a cubic polynomial in Weierstrass form. To this end, first define the mapping $f : (x, y) \mapsto \left(x, \frac{y+Cx+F}{2}\right)$. A tedious computation shows that φ transforms the affine part of X' to the curve X'' given by $\frac{y^2-F^2-C^2x^2}{4} - \frac{CFx}{2} = x^3 + Bx^2 + Ex + G$, i.e.,

$$y^2 = 4x^3 + (C^2 + 4B)x^2 + (4E + 2CF)x + (4G + F^2).$$

Substituting $2y$ for y , we can write X'' as

$$y^2 = x^3 + \underbrace{\left(\frac{C^2}{4} + B\right)}_{k_1} x^2 + \underbrace{\left(E + \frac{CF}{2}\right)}_{k_2} x + \underbrace{\left(G + \frac{F^2}{4}\right)}_{k_3}.$$

Next, define the mapping $g : (x, y) \mapsto \left(x - \frac{k_1}{3}, y\right)$. By another tedious computation, we see that g sends X'' to the curve given by

$$y^2 = x^3 + \left(k_2 - \frac{(k_1)^2}{3}\right)x + \left(k_3 + \frac{2(k_1)^3}{27} - \frac{k_1 k_2}{3}\right).$$

Since f and g are invertible affine maps, we are done.

2. Note that any linear change of variables preserving Weierstrass form must look like $(x, y) \mapsto (r^2x, r^3y)$ with $r \neq 0$. Let $J = \frac{A^3}{B^2}$. It's straightforward to check that J is invariant under any affine map of the form $(x, y) \mapsto (r^2x, r^3y)$.
3. Let $j = (27)(4^4) \left(\frac{A^3}{4A^3+27B^2}\right)$. This quantity is well-known as the *j-invariant*. Though tedious, checking that j is invariant under any suitable linear change of variables is still straightforward. In addition, elementary algebra shows that $J = \frac{j}{(27)(4^4)} + \frac{A^3(4A^3+27B^2)}{B^2(4A^3+27B^2)}$. ■

2 Complex analysis

2.1 Lecture 5

First, let's review some basic concepts about functions of a single complex variable.

Definition 2.1.1. Let $z_0 \in \mathbb{C}$. A function $f = u + iv : U \subset \mathbb{C} \rightarrow \mathbb{C}$ is *holomorphic* or *analytic* if at least one of the following equivalent conditions holds.

- (i) Both u and v are C^1 , and f satisfies the Cauchy-Riemann equations, i.e.,

$$\begin{aligned} u_x &= v_y \\ u_y &= -v_x. \end{aligned}$$

- (ii) $\frac{\partial f}{\partial \bar{z}} = 0$, where $\frac{\partial}{\partial \bar{z}} := \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$.

(iii) The Cauchy integral formula holds, i.e.,

$$f(w) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\eta)}{\eta - w} d\eta$$

for any closed circular path γ centered at w in U .

(iv) f has a power series representation on U .

Definition 2.1.2. A bijective function $f : U \subset \mathbb{C} \rightarrow V \subset \mathbb{C}$ is *biholomorphic* if it is holomorphic and its inverse is holomorphic. In this case, we say that U is *biholomorphic to* V , written as $U \approx V$.

Fact 2.1.3.

- (a) **Maximum modulus principle.** If $U \subset \mathbb{C}$ is a domain, $f : U \rightarrow \mathbb{C}$ is holomorphic, and $|f|$ has a local maximum, then f is constant.
- (b) **Liouville's theorem.** Any bounded entire function is constant.
- (c) **Riemann extension theorem.** If $\epsilon > 0$ and $f : B_{\epsilon}(z) \setminus \{z\} \subset \mathbb{C} \rightarrow \mathbb{C}$ is bounded and holomorphic, then f can be extended to a holomorphic function on $B_{\epsilon}(z)$.
- (d) **Riemann mapping theorem.** If $U \subsetneq \mathbb{C}$ is simply connected and open, then $U \approx B_1(0)$.
- (e) **Residue theorem.** If $f : B_{\epsilon}(0) \setminus \{0\}$ is holomorphic, then f can be expanded in a Laurent series $\sum_{n=-\infty}^{\infty} a_n z^n$ such that $a_{-1} = \frac{1}{2\pi i} \oint f(z) dz$.

Next, let's look at some basic concepts about functions of several complex variables.

Definition 2.1.4. A function $f = u + iv : U \subset \mathbb{C}^n \rightarrow \mathbb{C}$ is *holomorphic* if at least one of the following equivalent conditions holds.

- (i) f is holomorphic in each variable individually.
- (ii) Both u and v are C^1 , and f satisfies the Cauchy-Riemann equations,

$$\begin{aligned} u_{x_i} &= v_{y_i} \\ u_{y_i} &= -v_{x_i} \end{aligned}$$

for each $i = 1, \dots, n$.

(iii) $\sum_{i=1}^n \frac{\partial f}{\partial \bar{z}_i} = 0$.

(iv) f has a power series representation on U ,

$$\sum_{k_1=0}^{\infty} \cdots \sum_{k_n=0}^{\infty} a_{k_1, \dots, k_n} z_1^{k_1} \cdots z_n^{k_n}.$$

Note 2.1.5. Statements (a), (b), and (c) of Fact 2.1.3 generalize to functions of several variables, as does the Cauchy integral formula:

$$f(z) = \frac{1}{(2\pi i)^n} \int_{|\eta_i - z_i| = \epsilon_i} \frac{f(\eta_1, \dots, \eta_n)}{(\eta_1 - z_1) \cdots (\eta_n - z_n)} d\eta_1 \cdots d\eta_n$$

where $\eta_i > 0$ for each $i = 1, \dots, n$.

Theorem 2.1.6 (Hartog). *If $n > 1$, then any holomorphic function $f : B_\epsilon(0) \setminus \{0\} \subset \mathbb{C}^n \rightarrow \mathbb{C}$ extends to a holomorphic function on $B_\epsilon(0)$.*

Definition 2.1.7. Let X be a (topological) space. A sheaf F on X is a presheaf on X such that for any open $U \subset X$ and any open cover $\{U_i\}_{i \in J}$ of U , there is an exact sequence

$$0 \longrightarrow F(U) \longrightarrow F(U_i) \longrightarrow F(U_{ij})$$

where $U_{ij} := U_i \cap U_j$.

Example 2.1.8. Let S be a suitable object (such as a set or R -module) and $x \in X$. The *skyscraper sheaf* of X supported at x with value S is given by

$$\left(\begin{array}{c} U \subset X \\ \text{open} \end{array} \right) \mapsto \begin{cases} S & x \in U \\ \text{pt} & x \notin U \end{cases}.$$

Definition 2.1.9. A *ringed space* is a pair (X, \mathcal{J}) where X is a space and \mathcal{J} is a sheaf of rings on X .

Remark 2.1.10. Given any standard object (X, \mathcal{J}) , we can define a *geometric object* as a ringed space locally isomorphic to (X, \mathcal{J}) .

Definition 2.1.11 (Vector bundle). Let X and V be complex manifolds. Let $\pi : V \rightarrow X$ be holomorphic. We say that π is a (*holomorphic*) *vector bundle of rank n* if for any $x \in X$, there exist an open set $U \ni x$ in X and an isomorphism $\pi^{-1}(U) \xrightarrow{\cong} U \times \mathbb{C}^n$ such that the *transition maps* $U_{ij} \times \mathbb{C}^n \rightarrow U_{ij} \times \mathbb{C}^n$ are holomorphic and fiber linear.

Any vector bundle $\pi : V \rightarrow X$ induces a sheaf on X given by

$$F(U) = \Gamma(U, \pi^{-1}(U)).$$

Example 2.1.12.

1. The sheaf induced by the trivial bundle $\mathbf{1} := X \times \mathbb{C}$ is denoted by \mathcal{O}_X .
2. The tangent bundle TX of a smooth manifold X induces the sheaf of vector fields on X .
3. The cotangent bundle T^*X induces the sheaf $\Omega^1(X)$ of one-forms on X .
4. The alternating bundle $\wedge^p X$ of rank p induces the sheaf $\Omega^p(X)$ of p -forms on X .

3 Line bundles

3.1 Lecture 6

Definition 3.1.1. A *line bundle* is a vector bundle of rank 1.

Definition 3.1.2. Let X be a complex manifold. A *sheaf F of \mathcal{O}_X -modules* is a sheaf on X such that for any open set U in X ,

- (i) $F(U)$ is a module over $\mathcal{O}_X(U)$ and
- (ii) if $U \subset V \subset X$, then $(f \cdot a) \upharpoonright_U = f \upharpoonright_U \cdot a \upharpoonright_U$.

Example 3.1.3 (Sheaf of sections). Let X be a complex manifold and J be a vector bundle over X . For any open $U \subset X$, let

$$\mathcal{L}_J(U) = \Gamma(U, L).$$

This inherits a vector space structure from the family of fibers of V . Also, any relation of the form $U_1 \subset U_2 \subset U$ induces a linear map $\Gamma(U_2, L) \rightarrow \Gamma(U_1, L)$ given by $\sigma \mapsto \sigma \upharpoonright_{U_1}$. Thus, $\mathcal{L}_J(-)$ is a sheaf of vector spaces. Moreover, it is easily seen to be a sheaf of \mathcal{O}_X -modules.

Since any vector bundle is locally trivial, we see that \mathcal{L}_J is *locally free*, i.e., for any $x \in X$, there exist an (open) neighborhood U of x in X and an isomorphism $\varphi : \mathcal{L}_J(U) \rightarrow \bigoplus_{i=1}^{\text{rank}(J)} \mathcal{O}_X(U)$ such that for any open set $V \subset U$, the square

$$\begin{array}{ccc} \mathcal{L}_J(U) & \xrightarrow{\cong} & \bigoplus_{i=1}^{\text{rank}(J)} \mathcal{O}_X(U) \\ \downarrow & & \downarrow \\ \mathcal{L}_J(V) & \xrightarrow{\cong} & \bigoplus_{i=1}^{\text{rank}(J)} \mathcal{O}_X(V) \end{array}$$

commutes. In other words, \mathcal{L}_J is locally isomorphic to $(\mathcal{O}_X)^{\oplus \text{rank}(J)}$.

Definition 3.1.4. A sheaf F on a complex manifold X is *invertible* if there exist an open cover $\{U_i\}$ of X and a family of holomorphic isomorphisms $\varphi_i : \mathcal{O}_{U_i} \rightarrow \mathcal{L}_J \upharpoonright_{U_i}$.

Example 3.1.5. If J is a line bundle, then \mathcal{L}_J is invertible.

Consider the composite

$$\mathcal{O}_{U_i \cap U_j} \xrightarrow{\varphi_i} \mathcal{L}_J \upharpoonright_{U_i \cap U_j} \xrightarrow{\varphi_j^{-1}} \mathcal{O}_{U_i \cap U_j}, \quad 1 \mapsto g_{ij}.$$

From this, we can construct a line bundle L over X by defining the total space as

$$\coprod_i (U_i \times \mathbb{C}) / \sim$$

where $(x, \lambda)_i \sim (y, \mu)$ if $x = y$ and $\mu = g_{ij}\lambda$.

Definition 3.1.6 (Divisor). A *divisor on a complex manifold X* is a locally finite \mathbb{Z} -combination of irreducible holomorphic hypersurfaces of X . Equivalently, it is a subset of X locally defined by the vanishing of a holomorphic function.

Example 3.1.7. If $X = \mathcal{A}^1$, then any divisor D on X is of the form

$$D = \sum m_i p_i, \quad p_i \in \mathcal{A}^1, \quad m_i \in \mathbb{Z}.$$

Terminology. Each m_i is known as the *multiplicity of p_i* .

Any divisor D defines a line bundle $\mathcal{O}_X(D)$ on X and a holomorphic map $X \dashrightarrow \mathbb{P}(V^\vee)$ where $V \equiv \Gamma(X, \mathcal{O}_X(D))$. It is also true that any line bundle defines a divisor. It follows that

$$(\text{line bundles}) \xleftrightarrow{\sim} (\text{invertible sheaves}) \xleftrightarrow{\sim} (\text{divisors module linear equiv.}) . \quad (\dagger)$$

Consider the case where $D = \text{pt}$. Let $f \in \Gamma(U, \mathcal{O}_U)$ and let $U_i = X \setminus D$, which is a tubular neighborhood of D . Note that $U_i = f^{-1}(\mathbb{C} \setminus \text{hyperplane})$. Define $\mathcal{O}_X(D)$ as the line bundle with transition functions of the form $f|_{U_i \cap U_j}$.

Alternatively, let

$$(\mathcal{O}_X(D))(U) = \left\{ g : U \rightarrow \mathbb{C} \mid g \text{ is meromorphic, } \overbrace{fg}^{\text{product}} \text{ is holomorphic} \right\}.$$

For example, let $X = \mathbb{P}^1$ and D be a point p . Let (x_0, x_1) denote local coordinates on X near p . Let g be meromorphic in these coordinates and let $f(x_0, x_1) = \frac{x_1}{x_0}$. Then fg is holomorphic, i.e., g has a pole of order at most one at p .

Question.

1. What is $\Gamma(\mathbb{P}^1, \mathcal{O}_X)$?
2. What is $\Gamma(\mathbb{P}^1, \mathcal{O}_X(D))$?

In fact, it can be shown that

$$\Gamma(\mathbb{P}^1, \mathcal{O}_X(m, p)) = \begin{cases} \mathbb{C}\langle 1, x, \dots, x^m \rangle & m \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

In general, D is defined locally, and thus so is $\mathcal{O}_U(D)$. Specifically, $\Gamma(U, \mathcal{O}_U(D))$ consists of all holomorphic functions $f : U \setminus \text{supp}(D) \rightarrow \mathbb{C}$ such that if $D = \sum m_i Y_i$ and $Y_i \cap U = \{f_i = 0\}$, then $g \prod_i f_i^{m_i}$ is holomorphic in U .

Example 3.1.8 (Veronese embedding). Let $X = \mathbb{P}^1$ and p be as before.

1. Let $D = \mathcal{O}(2p)$. Consider the space $V := \Gamma(\mathbb{P}^1, \mathcal{O}(2p)) = \mathbb{C}\langle 1, x, x^2 \rangle$. Define the map $\varphi_{\mathcal{O}(2p)} : \mathbb{P}^1 \rightarrow \mathbb{P}^2$ by

$$\varphi_{\mathcal{O}(2p)}(x) = \underbrace{(1, x, x^2)}_{(x, y, z)}.$$

This is an embedding. Its image is precisely the smooth curve given by $y^2 = xz$.

2. Let $D = \mathcal{O}(3p)$. Then the image of the map $\varphi_{\mathcal{O}(3p)} : \mathbb{P}^1 \hookrightarrow \mathbb{P}^3$ given by $x \mapsto (1, x, x^2, x^3)$ is a so-called twisted cubic.

The line bundle L on X determines the map $X \dashrightarrow \mathbb{P}(\Gamma(X, L)^\vee)$ directly, as follows.

$$x \mapsto \ker \left(\Gamma(X, L) \xrightarrow{\text{eval}_x} L_p \right)$$

Definition 3.1.9. The *base locus* of L is $\mathcal{BL}(L) \equiv \{x \in X \mid s(x) = 0 \text{ for each } s \in \Gamma(X, L)\}$.

Note that we get a map $X \setminus \mathcal{BL}(L) \rightarrow \mathbb{P}(\Gamma(X, L)^\vee)$.

Now, let's consider a slight generalization of our preceding discussion. Let $V \subset \Gamma(X, L)$. This induces a map

$$\begin{array}{ccc} X & \dashrightarrow & \mathbb{P}(V^\vee) \\ \uparrow & \nearrow & \\ X \setminus \mathcal{BL}(V) & & \end{array} .$$

Let $X = \mathbb{P}^1$ and $p = \{x = 0\}$. Then $V \subset \Gamma(\mathbb{P}^1, \mathcal{O}(2)) = \mathbb{C}\langle 1, x, x^2 \rangle$, and

$$\begin{array}{ccc} \mathbb{P}^1 & \xrightarrow{\varphi_{\mathcal{O}(2)}} & \mathbb{P}^2 \\ & \searrow \varphi_V & \downarrow \rho \\ & & \mathbb{P}^1 \end{array}$$

commutes where ρ denotes the linear projection. Note that φ_V is a morphism so long as the center of ρ is not in the image of $\varphi_{\mathcal{O}(2)}$. In this case, we have that

$$\begin{aligned} \varphi_{\mathcal{O}(2)}(x) &= \frac{a + by + cx^2}{d + ex + fx^2} \\ \rho(x) &= \frac{a + bx}{c + dx}. \end{aligned}$$

3.2 Lecture 7

Let L_1 and L_2 be line bundles over X with transition functions $\{g_1^{kl} : U_{kl} \rightarrow \mathbb{C}^*\}$ and $\{g_2^{ij} : U_{ij} \rightarrow \mathbb{C}^*\}$, respectively. We can take a refinement $\{U_i \cap U_k\}$ where both L_1 and L_2 are trivial. Define $L^1 \otimes L^2$ as the line bundle with transition functions $\{g_1^{kl} g_2^{ij} : U_{ij} \cap U_{kl} \rightarrow \mathbb{C}^*\}$. Further, define $(L^1)^{-1}$ as the line bundle with transition functions $\{(g_1^{kl})^{-1} : U_{kl} \rightarrow \mathbb{C}^*\}$. Note that, locally, $L^1 \otimes (L^1)^{-1} \cong \mathcal{O}_X$.

Definition 3.2.1. We say that a divisor $D = \sum_i m_i Y_i$ is effective if $m_i \geq 0$ for each i .

Let $V = \Gamma(X, \mathcal{O}_X(D))$ and let D be effective. Note that $\mathbb{C}\langle D \rangle \subset V$. We have that $\text{supp}(D) = \varphi^{-1}$ (hyperplane) where $(\mathbb{C}\langle 0 \rangle)^\perp$ is precisely the hyperplane in $\mathbb{P}(V^\vee)$.

Example 3.2.2. Let $X = \mathbb{P}^1$.

1. Let $x = \frac{x_1}{x_0}$ and $D = p := \{x = 0\}$. Then $V = \mathbb{C}\langle 1, x \rangle$, and the map $\varphi_V : \mathbb{P}^1 \rightarrow \mathbb{P}(V^\vee)$ is given by $c \mapsto y := \frac{x}{1}$.
2. Let $D = m(\infty)$ with $m > 0$. Then $V = \mathbb{C}\langle x_1, \dots, x_m^m \rangle$, and the map $\varphi_{m\infty} : \mathbb{P}^1 \rightarrow \mathbb{P}^m$ is given by

$$\begin{aligned} (x_0, x_1) &\mapsto (x_0^m, x_0^{m-1}x_1, \dots, x_0x_1^{m-1}, x_1^m) \\ x &\mapsto (1, x, \dots, x^m). \end{aligned}$$

3. Let $D = p_1 + \dots + p_m$ where $p_i = [1 : t_i]$. Let $x = \frac{x_1}{x_0}$, so that ∞ is given by $x_0 = 0$. Then $V = \mathbb{C}\left\langle 1, \underbrace{\frac{1}{x-t_1}, \dots, \frac{1}{x-t_m}}_{a_0, \frac{a_1}{x-t_1}, \dots, \frac{a_m}{x-t_m}} \right\rangle$. This can be viewed as the space of all regular meromorphic functions on open subsets of \mathbb{P}^1 having poles of order at most m . The image of $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^m$ is precisely the hyperplane $\{a_0 = 0\}$.

Example 3.2.3. Let X be an elliptic curve, i.e., a space of the form \mathbb{C}/Λ . Let p be the image of 0 and let $D = mp$.

1. Let $m = 1$. Then $V = \Gamma(X, \mathcal{O}_X(D))$, which consists of all maps $f : X \rightarrow \mathbb{P}^1$ such that $f^{-1}(\infty) = \{0\}$. These are precisely the constant maps, so that $V \cong \mathbb{C}\langle s \rangle$ where s is a holomorphic section of $\mathcal{O}_X(D)$ vanishing at p and is meromorphic on \mathcal{O}_X .

$$\begin{array}{ccc} X & \dashrightarrow & \mathbb{P}^0 \\ \uparrow & \nearrow & \\ X \setminus p & & \end{array}$$

It follows that $\mathcal{BL}(\mathcal{O}_X(D)) = p$.

2. Let $m = 2$. Then $V = \mathbb{C}\langle 1, p \rangle$, and $\varphi_{2p} : X \rightarrow \mathbb{P}^1$ is precisely the D -th Weierstrass function. Moreover, we have an exact sequence

$$0 \longrightarrow \mathcal{O}_X(p) \longrightarrow \mathcal{O}_X(2p) \longrightarrow \mathcal{O}_p \longrightarrow \cdots$$

3. Let $m = 3$. Then $V = \langle 1, p, p' \rangle$, and the image of $\varphi_{3p} : X \rightarrow \mathbb{P}^2$ is given by $y^2 = x^3 + ax + b$.

Example 3.2.4. Let $X = \mathbb{P}^2$. Let $D = m \underbrace{(\text{line at } \infty)}_{\{z=0\}}$.

1. Let $m = 0$. Then $V = \mathbb{C}\langle 1 \rangle$, and $\mathcal{BL} = \emptyset$.
2. Let $m = 1$. Then $C = \mathbb{C}\langle \frac{x}{z}, \frac{y}{z}, \frac{z}{z} \rangle \cong \mathbb{C}\langle 1, X, Y \rangle$, and $\mathcal{BL} = \emptyset$. The map $\varphi_D : \mathbb{P}^2 \rightarrow \mathbb{P}^2$ is precisely the identity.
3. Let $m = 2$. Then $V = \langle \frac{x^2}{z^2}, \frac{x^4}{z^2}, \frac{y^2}{z^2} \cdot \frac{x}{z}, \frac{y}{z}, \frac{z}{z} \rangle$, and the map $\varphi_D : \mathbb{P}^2 \rightarrow \mathbb{P}^5$ is an embedding given by $(x, y, z) \mapsto \langle x^2, xy, y^2, xz, yz, z^2 \rangle$.

In general, if $H \subset \mathbb{P}^n$ is a hyperplane, then $\varphi_{\mathcal{O}(dH)} : \mathbb{P}^n \rightarrow \mathbb{P}^{\binom{d+n}{n}-1}$ is given by

$$(x_0, \dots, x_n) \mapsto (d\text{-th order homogenous polynomials}),$$

known as the d -th order Veronese embedding on \mathbb{P}^n .

Example 3.2.5. Let $X = \mathbb{P}^2$ with coordinates (x, y, z) . Let H denote the hyperplane given by $z = 0$ and let $D = 2H$. Then $V = \{s \in \Gamma(\mathcal{O}(2H)) \mid s(0, 0, 1) = 0\}$, and

$$\begin{array}{ccc} V & \hookrightarrow & \Gamma(\mathcal{O}(2H)) \\ \cong \uparrow & & \uparrow \cong \\ \mathbb{C}\langle x^2, xy, y^2, xz, yz \rangle & \hookrightarrow & \mathbb{C}\langle x^2, xy, y^2, xz, yz, z^2 \rangle \end{array}$$

commutes. Further, $\mathcal{BL}(V) = \{0\} = [0, 0, 1]$, and φ_V is a map $\mathbb{P}^2 \setminus \{0\} \rightarrow \mathbb{P}^4$ but does not extend to \mathbb{P}^2 . Indeed, we have that

$$\begin{aligned} \lim_{\substack{(0,y,1) \\ y \rightarrow 0}} \varphi_V &= \lim_{y \rightarrow 0} (0, 0, y^2, 0, y) = (0, 0, 0, 0, 1) \\ &\quad \Downarrow \\ \lim_{\substack{(x,0,1) \\ x \rightarrow 0}} \varphi_V &= \lim_{x \rightarrow 0} (x^2, 0, 0, x, 0) = (0, 0, 0, 1, 0). \end{aligned}$$

Note that for any $p \in X$, there exist \tilde{X} and $\pi : \tilde{X} \rightarrow X$ such that π restricted to $\pi^{-1}(X \setminus p)$ is an isomorphism and $\pi^{-1}(p)$ is a divisor on \tilde{X} that is isomorphic to \mathbb{P}^1 .

Proposition 3.2.6. *Let $Y \subset X$ be a submanifold of codimension $k \geq 2$. Let $\varphi : X \setminus Y \rightarrow Z$. Then there exist \tilde{X} and $\pi : \tilde{X} \rightarrow X$ such that π restricted to $\pi^{-1}(X \setminus Y)$ is an isomorphism and restricted to $\underbrace{\pi^{-1}(Y)}_{\text{divisor on } X}$ is a bundle with each fiber isomorphic to \mathbb{P}^{k-1} .*

Notation. In this case, the space \tilde{X} is denoted by $\text{Bl}_Y(X)$.

3.3 Lecture 8

Recall our correspondence (†). We can add to it the class of all maps

$$X \setminus \mathcal{BL}(L) \rightarrow \mathbb{P}(\Gamma(X, L)^\vee).$$

Let's turn now to some higher-dimensional examples.

Example 3.3.1. Let $X = \mathbb{P}^2$, $L = \mathcal{O}(2)$, and $V = \{s \in \Gamma(X, \mathcal{O}(2)) \mid \text{linearity condition}\}$. Then $\varphi_V : \mathbb{P}^2 \hookrightarrow \mathbb{P}^5$. Consider any homogenous polynomial $\sum a_{ijk}x^i y^j z^k$. Then our linearity condition may take any of the following forms.

- $\sum a_{ijk}x^i y^j z^k = 0$ where a_{ijk} ranges over

$$\{a_{000}, a_{120}, a_{020}, a_{101}, a_{011}, a_{002}\}.$$

- $a_{002} = 0$
- $a_{002} + a_{001} = 0$.

In the case of either of these last two, we get a map

$$\mathbb{P}^2 \xrightarrow{\varphi_V} \mathbb{P}^5 \xrightarrow{\psi} \mathbb{P}^4$$

for any $p \in \mathbb{P}^5$. There are two scenarios to consider.

- Suppose that $p \notin \text{im } \varphi_V$. Then $\psi \circ \varphi_V$ is a morphism.
- Suppose that $p = \varphi_V(001)$. Then ψ blows up at p . Consider the map $\varphi_V : \mathbb{P}^2 \setminus p \hookrightarrow \mathbb{P}^4$ given by $(x, y, z) \mapsto \underbrace{(x^2, xy, y^2, xz, y^2)}_{(x, y, z, u, v)}$. The image of this map is precisely $\text{im } \varphi_V \coprod_{\{x=y=z=0\}} \mathbb{P}^1 \subset \mathbb{P}^4$.

Terminology. In this setting, \mathbb{P}^1 is called an *exceptional divisor*.

Note that the equations

$$\begin{aligned} xz &= y^2 \\ zu &= yv \\ xv &= yu \end{aligned}$$

together generate the relevant ideal.

Remark 3.3.2. If we took L to be $\mathcal{O}(n)$ with $n \neq 2$, then our generators would still be quadratic.

Now, fix a and b and let $x = \epsilon a$, $y = \epsilon b$, and $z = 1$ where $\epsilon \rightarrow 0$. Then

$$\begin{aligned}\varphi_V(x, y, z) &= (\epsilon^2 a^2, \epsilon^2 ab, \epsilon^2 b^2, \epsilon a, \epsilon b) \\ &\sim (\epsilon a^2, \epsilon ab, \epsilon b^2, a, b) \\ &\rightarrow (0, 0, 0, a, b).\end{aligned}$$

Question. Is $\text{im } \varphi_V$ a manifold at $00010 = \varphi_V(1, b, a)$?

We have that

$$\begin{aligned}zu - yv &\rightarrow \frac{z}{u} = \frac{y}{u} \frac{v}{u} \\ xv = yu &\rightarrow \frac{x}{u} \frac{v}{u} = \frac{y}{u}.\end{aligned}$$

More generally, let X be a complex n -manifold and let $p \in X$. Then $\text{Bl}_p X = (X \setminus p) \amalg \underbrace{\mathbb{P}^{n-1}}_{\mathbb{P}(T_p X)}$. There at least two ways of extending the map

$$X \setminus p \xrightarrow{\varphi_V} \mathbb{P}^n \dashrightarrow^{\psi} \mathbb{P}^{n-1}$$

so that its image is a manifold at every point.

- (a) Provided that $\psi \circ \varphi_V$ is an embedding, then we can take $\text{Bl}_p(X)$ to be the closure of $X \setminus p$ in \mathbb{P}^{n-1} .
- (b) Let U is any polydisk containing the origin. We can replace $(X \setminus p) \cup U$ with $(X \setminus p) \cup \tilde{U}$ where \tilde{U} denotes the blow-up of U at 0.

More generally still, let $Y^m \subset X^n$ be a closed submanifold. Then $\tilde{X} := \text{Bl}_Y(X) = (X \setminus Y) \amalg \underbrace{\mathbb{P}(N_Y X)}_{\text{normal bundle}}$.

$$\begin{array}{ccc}\mathbb{P}^{n-m-1} & \longrightarrow & \mathbb{P}(N_Y X) \\ & & \downarrow \\ & & Y\end{array}$$


We wish to find a line bundle L over Y and a subspace $V \subset \Gamma(X, L)$ such that $\mathcal{BL}_V = Y$. In this case, the closure of the image of $\varphi_V : X \setminus Y \rightarrow \mathbb{P}(V^\vee)$ determines $(X \setminus Y) \cup \tilde{U}$ on U where U denotes any tubular neighborhood of Y in X .

Alternatively, if we are given an embedding

$$\begin{array}{ccc}Y & \hookrightarrow & X \\ \downarrow & & \downarrow \\ \mathbb{P}^{n-c} & \hookrightarrow & \mathbb{P}^n\end{array}$$

where c denotes the codimension of Y , then we can take $\text{Bl}_Y(X)$ to be the closure of $\text{Bl}_{\mathbb{P}^{n-c}}(\mathbb{P}^n \cap (X \setminus Y))$.

Example 3.3.3. Consider \mathbb{P}^3 with coordinates (x, y, z, w) . We wish to resolve the cone $\{x^2 = y^2\} \subset \mathbb{P}^3$. Let $p = \{x = z = 0\}$. We have a commutative diagram




Then the exceptional divisor in $\text{Bl}_p(\mathbb{P}^2)$ is isomorphic to $\mathbb{P}^2 \cong \mathbb{P}(T_p\mathbb{P}^2)$, and the exceptional divisor in $\text{Bl}_p(X)$ is isomorphic to the cone.

Example 3.3.4. Consider the quadratic map $\varphi : \mathbb{P}^2 \rightarrow \mathbb{P}^2$ given by $(x, y, z) \mapsto \left(\frac{1}{x}, \frac{1}{y}, \frac{1}{z}\right) = \underbrace{(yz, xz, xy)}_{(u,v,w)}$. Let

$$V = \{s \in \Gamma(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2)) \mid s(001) = 0, s(010) = 0, s(100) = 0\},$$

which is isomorphic to $\Gamma(\underbrace{\mathcal{I}_{3 \text{ points}}}_{\text{ideal sheaf on 3 points}} \otimes \mathcal{O}(2))$. The fact that $\varphi^{-1} = \varphi$ yields the following properties.

- The line $z = 0$ collapses to the point $u = v = 0$.
- The line $y = 0$ collapses to the point $u = v = 0$.
- The point $y = z = 0$ blows up to the line $u = 0$.



This hexagon is called the *del Pazzo surface of degree three*, denoted by dP_3 . Each of its lines is isomorphic to \mathbb{P}^1 .

Note 3.3.5. Suppose that C is a smooth curve and that $\dim X < 2$. Then $\varphi : C \setminus \text{pt} \rightarrow X$ automatically extends. But if C were singular or $\dim X \geq 2$, then this would be false.

3.4 Lecture 9

Definition 3.4.1 (Picard group). Let X be a complex manifold. The *Picard group* $\text{Pic}(X)$ of X is the group of all isomorphism classes of line bundles over X under \otimes .

Let $n \in \mathbb{N}$ and consider the family of line bundles $\{\mathcal{O}(k) \mid k \in \mathbb{Z}\}$ over \mathbb{P}^n .

Proposition 3.4.2. $\text{Pic}(\mathbb{P}^n) \cong \mathbb{Z}$ with generator $\mathcal{O}(1)$.

Let $\mathbb{P}^n = \mathbb{P}(V)$. We have an exact sequence

$$0 \longrightarrow L \longrightarrow \mathcal{O}_{\mathbb{P}^n} \otimes_{\mathbb{C}} V \longrightarrow \dots$$

We have that

1. $\Gamma(\mathbb{P}^n, \mathcal{O}(1)) = \mathbb{C}\langle z_0, \dots, z_n \rangle = V^\vee$,
2. $\Gamma(\mathbb{P}^n, \mathcal{O}(-1)) = 0$, and
3. $\Gamma(\mathbb{P}^n, \mathcal{O}(k)) = \begin{cases} \text{Sym}^k(V^\vee) & k \geq 0 \\ 0 & k < 0 \end{cases}$.

Let $U_i = \{z \in \mathbb{P}^n \mid z_i \neq 0\}$ for each $i \in \{0, 1, \dots, n\}$, so that $\mathbb{P}^n = \bigcup_{i=0}^n U_i$. Let $Z_{ij} = \frac{z_j}{z_i}$, thereby endowing each U_i with local coordinates. Let s be a section of \mathcal{O} , so that

$$s = (s_i \in \Gamma(U_i, \mathcal{O}))_{i=0}^n.$$

Note that Z_i defines a section on U_j with $s_j = \frac{z_i}{z_j} = Z_{ji}$ for each $j = 0, \dots, n$.

$$\begin{array}{ccc} s_j & \longleftarrow & Z_{jk} \cdot s_k \\ \parallel & & \parallel \\ \frac{z_i}{z_j} & \longleftarrow & Z_{jk} \cdot \frac{z_i}{z_k} \end{array}$$

We can establish the following properties.

1. If $\mathcal{O} = \mathcal{O}(1)$, then $s_i = Z_{ij}s_j$.
2. If $\mathcal{O} = \mathcal{O}(-1)$, then $s_i = Z_{ji}s_j$.
3. If $\mathcal{O} = \mathcal{O}(k)$, then $s_i = (Z_{ij})^k s_j$.

In summary,

	\mathcal{O} (trivial)	$\mathcal{O}(-1)$	$\mathcal{O}(1)$	$\mathcal{O}(k)$
LB	$\mathbb{P}^n \times \mathbb{C}$	tautological	dual	
Sheaf	1	Z_{ji}	Z_{ij}	$(Z_{ij})^k$
Divisor	0	$-H$ h.p.	$+H$	kH
Map	pt	undefined	id	$\left\{ \begin{array}{ll} \text{Veronese} & k > 0 \\ \text{undefined} & k < 0 \\ \text{pt} & k = 0 \end{array} \right.$

Let X be a complex n -manifold. Then T_X consists of all local sections on an open set U with coordinates, say, z_1, \dots, z_n . The set $\left\{ \frac{\partial}{\partial z_i} \right\}$ is a basis for this, with each section of the form $\sum_{i=1}^n f_i \frac{\partial}{\partial z_i}$ where each f_i belongs to $\Gamma(U, \mathcal{O})$. For any other basis $\left\{ \frac{\partial}{\partial w_i} \right\}$, we have that

$$\frac{\partial}{\partial w_i} = \sum \frac{\partial z_j}{\partial w_i} \frac{\partial}{\partial z_j}.$$

Note that $T_V \cong \mathcal{O}_V \otimes_{\mathbb{C}} V$. In general, $\Omega_V^i \cong \mathcal{O}_V \otimes \wedge^i V^\vee$.

Question. What is $T_{\mathbb{P}(V)}$?

Note 3.4.3 (Bundle associated to an n -manifold).

1. $T_X^\vee = \Omega \equiv \Omega^1$, whose transition functions are precisely the inverses of the transposes of those for T_X .
2. Let $\Omega^i = \wedge^i \Omega^1$. If $i = n$, then we call this space the *canonical sheaf* K_X or the *dualized sheaf* ω_X .
3. Recall the map $\wedge^i : \mathrm{GL}(n) \rightarrow \mathrm{GL}\left(\binom{n}{i}\right)$. If $i = n$, then this is precisely the determinant map.

Consider the exact sequence

$$0 \longrightarrow \mathcal{O} \longrightarrow \mathcal{O}(1)^{\oplus n+1} \longrightarrow T_{\mathbb{P}^n} \longrightarrow 0$$

$$1 \longmapsto (z_i)$$

$$(a_i) \longmapsto \sum a_i \frac{\partial}{\partial z_i}$$

Terminology. The vector field given by $\sum z_i \frac{\partial}{\partial z_i}$ is known as the *Euler vector field*.

Moreover, we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \underbrace{\mathcal{O}_{\mathbb{P}(V)}}_{\mathbb{C}} & \longrightarrow & \underbrace{\mathcal{O}_{\mathbb{P}(V)}(1) \otimes V}_{V^\vee} & \longrightarrow & T_{\mathbb{P}(V)} \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & \mathcal{O}_V(1) \otimes V & \xrightarrow{\cong} & T_V \longrightarrow 0 \end{array}$$

Terminology. The top row of this diagram is known as the *Euler sequence*.

Therefore, the *weight* of V equals -1 , whereas the weight of V^\vee equals $+1$.

Informally, any holomorphic function f on V is the same as a direct sum of homogenous functions of degree k , i.e., has the form

$$\bigoplus_{k=0}^{\infty} \Gamma(\mathbb{P}(V), \mathcal{O}(k)),$$

called the *Taylor expansion of f* .

Note 3.4.4. In general, we have an exact sequence

$$0 \longrightarrow \mathcal{O}(n) \longrightarrow \mathcal{O}^{(n+1)} \longrightarrow T_{\mathbb{P}}(-1) \longrightarrow 0,$$

which becomes the Euler sequence

$$0 \longrightarrow \mathcal{O} \longrightarrow \mathcal{O}(1) \oplus \mathcal{O}(1) \longrightarrow T_{\mathbb{P}^1} \longrightarrow 0$$

in the case where $n = 1$. It follows that

$$K_{\mathbb{P}^n} = \mathcal{O}_{\mathbb{P}^n}(-n-1).$$

Lemma 3.4.5. *If $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is an exact sequence of vector spaces, then*

$$\det(B) = \det(A) \otimes \det(C).$$

Corollary 3.4.6. $\mathcal{O}(2) \cong \det(\mathcal{O}(1) \oplus \mathcal{O}(1)) = \det(\mathcal{O}) \otimes \det(T) = \det(T)$.

Remark 3.4.7. Similarly, we can show that $\det(T_{\mathbb{P}^n}) = \mathcal{O}_{\mathbb{P}^n}(n+1)$.

Suppose that $X \subset Y$ is a submanifold of codimension 1. Then we have a short exact sequence

$$0 \longrightarrow T_X \longrightarrow (T_Y)|_X \longrightarrow N_{X/Y} \longrightarrow 0.$$

Lemma 3.4.8. $N_{X/Y} \cong \mathcal{O}_Y(X)|_X$.

In other words, if $L \in \text{Pic}(Y)$, $s \in \Gamma(Y, L)$, and $X = \{s = 0\}$, then $N_{X/Y} \cong L|_X$.

Theorem 3.4.9 (Adjunction formula). $K_X \cong (K_Y \otimes_{\mathcal{O}_Y} \mathcal{O}_Y(X))|_X$.

Proof. Note that $(K_Y^{-1})|_X = K_X^{-1} \otimes N_{X/Y}$. Thus,

$$\begin{aligned} K_X &\cong K_Y|_X \otimes N_{X/Y} \\ &\cong K_Y|_X \otimes \mathcal{O}_Y(X)|_X \\ &\cong (K_Y \otimes \mathcal{O}_Y(X))|_X. \end{aligned}$$

□

3.5 Lecture 10

Proof of Lemma 3.4.8. Let $s \in \Gamma(Y, L)$. We can write $s = fs_0$, so that $ds = s_0df + fds_0$. Consider the short exact sequence

$$0 \longrightarrow T_X \longrightarrow T_Y|_X \xrightarrow{ds} L \longrightarrow 0.$$

Thus, ds transforms just as s_0 does. □

Example 3.5.1.

1. Let $Y = \mathbb{P}^3$. Suppose that \tilde{X} is a smooth curve of degree d . Then $K_Y = \mathcal{O}(-3)$, and $K_X = \mathcal{O}(d-3)|_X$. Further, if g denotes the genus of a surface, then Bézout's theorem implies that

$$\begin{aligned} 2g - 2 &= \deg(K_X) = d(d-3) \\ &\quad \downarrow \\ g &= 1 + \frac{d(d-3)}{2} = \frac{(d-1)(d-2)}{2}. \end{aligned}$$

In particular,

d	g
1	0
2	0
3	1
4	3
5	6

2. Let $Y = \mathbb{P}^n$ and let $X \subset Y$ be of dimension d . Note that $K_X = \mathcal{O}_X$ precisely when $d = n + 1$. In particular,

n	X
2	cubic / elliptic curve
3	quartic (<i>a K_3 surface</i>)
4	quintic

Let $p_1, \dots, p_n \in \mathbb{P}^N$, let $m_1, \dots, m_n \in \mathbb{Z}_{\geq 1}$, and let $d \in \mathbb{Z}$. We wish to describe

$$\Gamma(\mathcal{J}_{\Sigma_{m_i p_i}}(d)) := (\mathcal{J}_{\Sigma_{m_i p_i}} \otimes \mathcal{O}(d)).$$

For simplicity, let $N = 2$.

Definition 3.5.2. If $n = 1$, then *imposition* is $\text{Imp}_m \equiv \text{codim}(\Gamma(\mathcal{J}_{mp}(d), \Gamma(\mathcal{O}(d))))$.

Proposition 3.5.3. $\text{Imp}_m = \binom{m+1}{2}$.

Definition 3.5.4. Consider the space Γ .

1. The *actual dimension* of Γ is the dimension of Γ as a vector space.
2. The *virtual dimension* $\text{vd}(\Gamma)$ of Γ is the quantity $\binom{d+2}{2} - 1 - \sum_i \binom{m_i+1}{2}$.
3. The *expected dimension* of Γ is the quantity $\max(\text{vd}(\Gamma), 0)$.

Conjecture 3.5.5. *The actual dimension always equals the expected dimension.*

Answer. This is **false**. For example, let $N = 2$, $d = 1$, $m_i = 1$, and $n = 3$. Then $\Gamma = 0$, so that $\mathbb{P}(\Gamma) = \emptyset$. Hence the expected dimension is zero, but the actual dimension is positive whenever the p_i are co-linear. \square

This leads us to the following modification of Conjecture 3.5.5.

Conjecture 3.5.6. *If the p_i are in general position, then the actual dimension equals the expected dimension.*

Answer. This is **false**. To see this, let $d = 2$ and $N = n = m_i = 2$. Consider a conic C through five points. Here, our conjecture holds. But if instead $N = 2$, $d = 4$, $n = 5$, and $m_i = 2$, then the virtual dimension is precisely $\binom{4+2}{2} - 5 \cdot 3 = 0$. Since the square of C exists, it follows that our conjecture fails. \square

We can improve Conjecture 3.5.6 as follows.

Conjecture 3.5.7. *If the actual dimension is different from the expected dimension, then $\Gamma(\mathcal{J}_{\sum m_i p_i}(d))$ has a base curve.*

Answer. This is **unknown**. See the article “[Linear Systems of Plane Curves](#)” by Rick Miranda. □

Consider the map $|\mathcal{O}(d)| : \mathbb{P}^2 \hookrightarrow \mathbb{P}^{\binom{d+2}{2}-1}$. We also have a map

$$\text{Bl}_{p_1, \dots, p_n}(\mathbb{P}^2) \xlongequal{\quad} \underbrace{\mathbb{P}^2}_{\widetilde{\mathbb{P}^2}} \begin{array}{c} \uparrow \\ \mathbb{P}^2 \end{array} \begin{array}{c} \xrightarrow{\mathcal{J}_{\sum p_i}(d)} \\ \mathbb{P}^{\dim-1} \end{array}$$

Proposition 3.5.8. *Consider the blow-up $\pi : \underbrace{\widetilde{\mathbb{P}^2}}_X \rightarrow \mathbb{P}^2$. We have that*

$$\text{Pic}(X) \cong \mathbb{Z} \langle \pi^*(\mathcal{O}(1)), E_1, \dots, E_n \rangle$$

where E_i denotes the divisor collapsing to p_i .

Remark 3.5.9.

Good: $\pi^*\mathcal{O}(d) - \sum m_i E_i \longleftrightarrow \mathcal{J}_{\sum m_i p_i}(d)$.

Better: $\Gamma(X, \cdot) = \Gamma(\mathbb{P}^2, \cdot)$.

Best: $\pi_*(\cdot) = \cdot$.

Conjecture 3.5.10. *Any line bundle $L := (\pi^*\mathcal{O}(d) - \sum m_i E_i)$ has the expected dimension of the space of sections unless $\mathcal{BL}(L)$ contains a (-1) -curve, i.e., a smooth curve C of genus zero such that $C^2 = -1$.*

Example 3.5.11 (*(-1) -curve*). Let $d = 1$, $n = 2$, and $m_1 = m_2 = 1$. If $C \in \mathcal{O}(1)(-p - q)$, then $C^2 = 1^2 - 1 - 1 = -1$. In general,

$$\mathcal{O}(d) \left(\left(- \sum m_i E_i \right) \left(\mathcal{O}(d) - \sum m_i p_i \right) \right) = dd' - \sum m_i m'_i.$$


In \mathbb{P}^2 , this means the number of intersections other than the p_i .

Space	C^2
$\mathcal{O}(1)$	1
$\mathcal{O}(1)(-p)$	0
$\mathcal{O}(1)(-p - q)$	-1
⋮	
$\mathcal{O}(2)$	4
$\mathcal{O}(2)(-p_1)$	3
$\mathcal{O}(2)(-p_1 - p_2)$	2
⋮	
$\mathcal{O}(2)(-p_1 - \dots - p_4)$	0
$\mathcal{O}(2)\left(-\sum_{i=1}^5 p_i\right)$	-1

4 Kähler manifolds

4.1 Lecture 11


Consider the following Hasse diagram of subgroups:



where $\text{Sp}(2n, \mathbb{R})$ denotes the group of real $2n \times 2n$ *symplectic matrices*, i.e., matrices M satisfying

$$M^t \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} M = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}.$$

Similarly, we can view various areas of geometry as refinements of certain others:



Before investigating Kähler geometry, let us establish some basic geometric concepts.

Definition 4.1.1. Let X be a real manifold. An *almost complex structure on X* is a bundle map $I : TX \rightarrow TX$ such that $I^2 = -1$.

Note that the eigenvalues of I are precisely i and $-i$.

Notation.

1. Let $T^{1,0}$ denote the eigenspace of i .
2. Let $T^{0,1}$ denote the eigenspace of $-i$.

Any complex manifold X has a natural almost complex structure. Indeed, given local coordinates x_i, y_i on X , define I by $\frac{\partial}{\partial x_i} \mapsto \frac{\partial}{\partial y_i}$ and $\frac{\partial}{\partial y_i} \mapsto -\frac{\partial}{\partial x_i}$. It follows that any manifold with an almost complex structure has even dimension.

Now, consider the complexification of our tangent bundle, $T^{\mathbb{C}}X \equiv TX \otimes_{\mathbb{R}} \mathbb{C}$.

Proposition 4.1.2.

1. $T^{\mathbb{C}}X \cong T^{1,0} \oplus T^{0,1}$.

$$2. T^{\mathbb{C}}X \cong T^{*1,0} \oplus T^{*0,1}$$

Define, formally, the complex coordinates $z_j = x_j + iy_j$. Note that $T^{\mathbb{C}}X$ has as basis $\left\{\frac{\partial}{\partial z_j}, \frac{\partial}{\partial \bar{z}_j}\right\}$ and that $T^{*\mathbb{C}}X$ has as basis $\{dz_j, d\bar{z}_j\}$ where $dz_j \equiv dx_j + idy_j$.

Notation.

1. $\bigwedge^k X := \bigwedge^k T^*X$.
2. $\bigwedge^{p,q} X := \bigwedge^p T^{*1,0}X \otimes_{\mathbb{C}} \bigwedge^q T^{*0,1}X$.

Note 4.1.3. Let X be an n -dimensional complex manifold.

1. $\left(\bigwedge^k T^*X\right) \otimes \mathbb{C} = \bigwedge_{\mathbb{C}}^k (T^*X \otimes \mathbb{C})$.
2. $\left(\bigwedge^k X\right) \otimes \mathbb{C} = \bigoplus_{p+q=k} \bigwedge^{p,q} X$.

Therefore, $\left(\bigwedge^k X\right) \otimes \mathbb{C}$ can be decomposed according to the counting equation $\binom{2n}{k} = \sum \binom{n}{p} \binom{n}{q}$.

Let U and V be open in \mathbb{C}^n . Let $f : U \rightarrow V$ be holomorphic. Then the map $df : TU \rightarrow TV$ extends to a map $df^{\mathbb{C}} : T^{\mathbb{C}}U \rightarrow T^{\mathbb{C}}V$ that preserves both $T^{1,0}$ and $T^{0,1}$.

Let $\mathcal{A}^{p,q} = \Gamma(\bigwedge^{p,q})$, i.e., $\mathcal{A}^{p,q}(U) = \Gamma(U, \bigwedge^{p,q})$. Consider the exterior derivative $d : \mathcal{A}^k \rightarrow \mathcal{A}^{k+1}$. With π denoting the projection map, define the operators

$$\begin{aligned} \partial &= \pi^{p+1,q} \circ d \\ \bar{\partial} &= \pi^{p,q+1} \circ d \end{aligned}$$

on $\mathcal{A}^{p,q}$. Locally, we have that

$$df = \frac{\partial f}{\partial x_i} dx_i + \frac{\partial f}{\partial y_i} dy_i = \frac{\partial f}{\partial z_i} dz_i + \frac{\partial f}{\partial \bar{z}_i} d\bar{z}_i$$

for any $f \in \mathcal{A}^{0,0}$. By the Cauchy-Riemann equations, it follows that f is holomorphic if and only if $\bar{\partial}f = 0$.

Remark 4.1.4. Any (p, q) -form locally looks like $f_{IJ} dz_I \wedge \bar{z}_J$.

Proposition 4.1.5.

1. $d = \partial + \bar{\partial}$.
2. $\partial^2 = 0 = \bar{\partial}^2$.
3. $\partial \bar{\partial} = -\bar{\partial} \partial$.
4. $\partial(\alpha \wedge \beta) = \partial\alpha \wedge \beta + (-1)^{p+q} \alpha \wedge \partial\beta$ for any $\alpha \in \mathcal{A}^{p,q}$ and $\beta \in \mathcal{A}^{r,s}$.

Lemma 4.1.6 (Single-variable Poincaré). Consider the disk $B_\epsilon \subset \overline{B_\epsilon} \subset U \subset \mathbb{C}$ where U is open. Let $\alpha = f d\bar{z} \in \mathcal{A}^{0,1}(U)$ and

$$g(z) = \frac{1}{2\pi i} \int_{B_\epsilon} \frac{f(w)}{w - z} dw \wedge d\bar{w}.$$

Then $\bar{\partial}g = \alpha$.

Lemma 4.1.7 (Multi-variable Poincaré). *Consider the polydisk $B_\epsilon \subset \overline{B_\epsilon} \subset U \subset \mathbb{C}^n$ where U is open. Let $\alpha \in \mathcal{A}^{p,q}$ with $q > 0$ and $\bar{\partial}\alpha = 0$. Then there is some $\beta \in \mathcal{A}^{p,q-1}(B_\epsilon)$ such that $\bar{\partial}\beta = \alpha$.*

Remark 4.1.8. If U is contractible, then any differential form on U is closed if and only if it is exact.

Let $U \subset \mathbb{C}^n$ be open and let I denote the natural almost complex structure on U . Let g be a Riemannian metric on U .

Definition 4.1.9 (Hermitian metric).

1. We say that g is *compatible with I* or (almost) *Hermitian* if $g(u, v) = g(Iu, Iv)$.
2. If g is Hermitian, then the real $(1, 1)$ -form $\omega \in \mathcal{A}^{1,1}(U) \cap \mathcal{A}^2(U)$ defined by

$$\omega(u, v) = g(Iu, v)$$

is called the *fundamental form of g* .

Notation. $h := g - i\omega$.

Definition 4.1.10. A Hermitian matrix M is *positive-definite* if $z^*Mz > 0$ for every nonzero complex column vector z .

Note that h is a positive-definite form in the sense that, locally, its component functions define a positive-definite matrix at any given point.

Example 4.1.11. Let $g = \underbrace{dx^2}_{dx \otimes dx} + dy^2 = \sum_{i=1}^n dx_i^2 + dy_i^2 \in T^* \otimes T^* \subset (T^* \otimes T^*) \otimes_{\mathbb{R}} \mathbb{C}$. Since

$$dz \wedge d\bar{z} = (dx + idy) \wedge (dx - idy) = -2idx \wedge dy,$$

it follows that

$$\omega = dx \otimes dy - dy \otimes dx = \frac{i}{2} dz \wedge d\bar{z}.$$

Moreover, we see that

$$\begin{aligned} h &= z - i\omega \\ &= dx^2 - idxdy + idydx + dy^2 \\ &= dx(dx - idy) + idy(dx + idy) \\ &= (dx + idy)(dx - idy) \\ &= dz \otimes d\bar{z}. \end{aligned}$$

For each $z \in \mathbb{C}^n$, define the matrix $(h_{ij})(z)$ by

$$h_{ij}(z_1, \dots, z_n) = h\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}\right).$$

Proposition 4.1.12. *Let I be an almost complex structure on $U \subset X$ and let g be compatible with I . Then $d\omega = 0$ if and only if for each $x \in X$, there exist a neighborhood U' of x and a holomorphic map $f : U' \rightarrow U$ such that f^*g oscillates the standard metric to the second order, i.e., $(h_{ij}) = \text{id} + O(|z|^2)$.*

Notation. In this case, we write $h \approx \text{id}$.

Definition 4.1.13 (Kähler manifold). Consider the four-tuple (X, I, g, ω) . We say that X is a *Kähler manifold* if $d\omega = 0$. In this case, we call g a *Kähler metric on X* and ω a *Kähler form*.

Definition 4.1.14. Let (X, I, g, ω) be a Kähler structure with $\dim X = n$.

1. The *Lefschetz operator* $L : \bigwedge^k X \rightarrow \bigwedge^{k+2} X$ is defined by $\alpha \mapsto \alpha \wedge \omega$.
2. The *Hodge $*$ -operator* $*$: $\bigwedge^k X \rightarrow \bigwedge^{2n-k} X$ is defined by the property

$$\alpha \wedge * \beta = \hat{g}(\alpha, \beta) \omega^n$$

where \hat{g} is induced by g and ω^n denotes the (positively oriented) volume form on X .

3. The *dual Lefschetz operator* $\Lambda : \bigwedge^k X \rightarrow \bigwedge^{k-2} X$ is defined as the composite $*^{-1} \circ L \circ *$.

Note 4.1.15.

1. In coordinates for which $h \approx \text{id}$, we have that $*dx^I = dx^{\partial}$ where $\partial := I^{\mathbb{C}}$??.
2. Λ is \mathcal{O} -linear.

4.2 Lecture 12

Proposition 4.2.1. *Let X be a complex manifold. Let ω be a closed real positive-definite form of type $(1, 1)$, i.e., locally, $\omega = \frac{i}{2} \sum h_{i\bar{j}} dz_i \wedge d\bar{z}_j$ such that the matrix $(h_{i\bar{j}}(p))$ is positive-definite for each p . Then there exists a Kähler metric g on X such that ω equals the fundamental form of g .*

Since every Kähler form is positive-definite, it follows that the set \mathbb{K}_X of all Kähler forms on X is precisely the set of all closed real positive-definite forms of type $(1, 1)$.

Definition 4.2.2. Let V be a vector space over \mathbb{R} . A subset $C \subset V$ is a *convex cone* if $av_1 + bv_2 \in C$ for any $v_1, v_2 \in C$ and any $a, b \in \mathbb{R}_{>0}$.

Corollary 4.2.3. *Suppose that X is compact. Then \mathbb{K}_X is an open convex cone in the infinite-dimensional real vector space $S := \{\omega \in \mathcal{A}^{(1,1)}(X) \cap \mathcal{A}^2(X) \mid d\omega = 0\}$.*

Idea. The fact that \mathbb{K}_X is a convex cone follows from the fact that the set of all positive-definite matrices is a convex cone. It remains to show that \mathbb{K}_X is open. Since X is compact, it has a finite open cover $\{U_i\}$. The set $P_{U_i} \subset S$ of all forms that are positive-definite on U_i is open. Thus, $\bigcap_i P_{U_i} = \mathbb{K}_X$ is also open. \square

Remark 4.2.4. It turns out that $S \cong H^2(X, \mathbb{R})$.

Example 4.2.5.

1. The form $\omega \equiv \frac{i}{2} dz \wedge d\bar{z}$ is Kähler on \mathbb{C} and is exact.
2. The same form descends to a Kähler form on the torus \mathbb{C}/Λ , which is not exact.
3. Consider the inclusion $i : X \rightarrow Y$ of a closed submanifold. If ω is Kähler on Y , then $i^*\omega$ is Kähler on X .

Note 4.2.6. Let $f : X \rightarrow Y$ be holomorphic and let ω be a Kähler form on Y . It is *not* necessarily true that $f^*\omega$ is Kähler on X . For example, if $f(x) = \mathbf{pt}$ for all $x \in X$, then $f^*\omega$ is the zero form and thus not positive. In general, f must be injective. For example, if $f : C \rightarrow \mathbb{C}$ is a double cover where C is a Riemann surface, then C inherits a Kähler form only outside the *ramification of f* , i.e., the set

$$\{c \in C \mid \text{there is no neighborhood } U \text{ of } c \text{ such that } f|_U \text{ is injective}\}.$$

This is precisely the set of points at which df is nonzero.

Example 4.2.7.

1. Consider the open cover $\{U_i\}_{1 \leq i \leq n}$ of \mathbb{P}^n where $U_i \equiv \{z \in \mathbb{P}^n \mid z_i \neq 0\}$. Define $\varphi_i : U_i \xrightarrow{\cong} \mathbb{C}^n$ by

$$(z_0, \dots, z_n) \mapsto \underbrace{\left(\frac{z_0}{z_i}, \dots, \frac{z_{i-1}}{z_i}, \frac{z_{i+1}}{z_i}, \dots, \frac{z_n}{z_i} \right)}_{(w_1, \dots, w_n)}.$$

Then $\{(U_i, \varphi_i)\}$ is a holomorphic atlas on \mathbb{P}^n . For each i , let

$$\omega_i = \frac{i}{2\pi} \partial \bar{\partial} \log \left(\sum_{l=0}^n \left| \frac{z_l}{z_i} \right|^2 \right).$$

By way of φ_i , this becomes

$$\frac{i}{2\pi} \partial \bar{\partial} \log \left(1 + \sum_{k=1}^n |w_k|^2 \right).$$

Exercise 4.2.8. Show that $\omega_i|_{U_i \cap U_j} = \omega_j|_{U_i \cap U_j}$.

Therefore, the ω_i patch together to form a metric ω on \mathbb{P}^n , known as the *Fubini-Study metric*.

Exercise 4.2.9. Show that ω is closed, real, positive, and of type $(1, 1)$.

It follows that ω is a Kähler metric.

2. Any branched cover of \mathbb{P}^n admits a Kähler metric (which must be different from the pullback of a Kähler metric on \mathbb{P}^n). For example, consider an elliptic curve $E \rightarrow \mathbb{P}^1$, which fits into a commutative square

$$\begin{array}{ccc} E & \longrightarrow & \mathbb{P}^1 \\ \parallel & & \uparrow \text{---} \\ E & \longleftarrow & \mathbb{P}^2 \end{array}.$$

Definition 4.2.10. A complex manifold is *projective* if it is isomorphic to a closed submanifold of projective space.

Proposition 4.2.11. Any projective complex manifold is Kähler.

Proof. This follows from Example 4.2.5(3) together with Example 4.2.7(1). \square

Definition 4.2.12. Let X be a complex manifold. Let D be a first-order operator on $\mathcal{A}^*(X)$.

1. The *adjoint* of D is

$$D^* \equiv -* \circ D \circ *$$

2. The *Laplacian associated to D* is

$$\Delta_D \equiv DD^* + D^*D.$$

Definition 4.2.13. The *Laplace operator* is $\Delta \equiv dd^* + d^*d$.

Example 4.2.14.

1. Let $D = \partial$. Then $\partial^* (f_{IJ} dz^I \wedge dz^J) = \sum_{i \in I} f_{IJ} dz^{I-i} \wedge dz^J$.

2. Let $D = d$. Let (x_1, \dots, x_n) be local coordinates on X . Then

$$\begin{aligned} d(f dx^I) &= \sum_{i \notin I} \frac{\partial f}{\partial x_i} dx^i \wedge dx^I \\ d^*(f dx^I) &= \sum_{i \in I} \frac{\partial f}{\partial x_i} dx^{I-i}. \end{aligned}$$

Therefore,

$$\begin{aligned} d \circ d^*(f dx^I) &= \frac{\partial^2}{\partial x_i \partial x_j} dx^{I-i \cup j} \\ &= \sum_{\substack{i \in I \\ j \notin I}} \dots + \sum_{i=j \in I} \dots \\ d^* \circ d(f dx^I) &= 0 + \sum_{i=j \notin I} \dots, \end{aligned}$$

so that $\Delta_D = \sum \frac{\partial^2 f}{\partial x_i^2}$.

Theorem 4.2.15 (Kähler identities). Let (X, I, g, ω) be a Kähler manifold.

1. $[\bar{\partial}, L] = 0 = [\partial, L]$.
2. $[\partial^*, \Lambda] = 0 = [\bar{\partial}^*, \Lambda]$.
3. $[\bar{\partial}^*, L] = i\partial$ and $[\partial^*, L] = -i\bar{\partial}$.
4. $\Delta_\partial = \Delta_{\bar{\partial}} = \frac{1}{2}\Delta$, and Δ commutes with $*$, ∂ , $\bar{\partial}$, ∂^* , $\bar{\partial}^*$, L , and Λ .

5 Lie algebras

A *Lie algebra* is a (finite-dimensional) vector space V endowed with a *bracket* $[-, -] : V \times V \rightarrow V$ satisfying bilinearity, antisymmetry, and the Jacobi identity.

A linear map $f : V \rightarrow W$ is a *Lie algebra homomorphism* if $f[x, y] = [f(x), f(y)]$ for all $x, y \in V$.

A (*Lie*) *ideal* of V is a subspace W of V such that $[W, V] \subset W$.

Definition 5.0.1. Let $(V, [-, -])$ be a Lie algebra.

1. We say that V is *abelian* if $[x, y] = 0$ for all $x, y \in V$.
2. We say that V is *simple* if it is non-abelian and has no non-trivial ideals.
3. We say that V is *semi-simple* if it is the direct sum of simple Lie algebras.
4. We say that v is *adjoint* if it is both semi-simple and centerless.

Now let G be any Lie group. For any $g \in G$, left-translation $\ell_g : G \rightarrow G$ is an isomorphism of \mathbb{C} -manifolds. Thus, if V is a vector field on G , then so is $(\ell_g)_* V$.

Definition 5.0.2.

1. We say that a vector field V on G is *left-invariant* if $(\ell_g)_* V = V$ for every $g \in G$.
2. The *Lie algebra* $\text{Lie}(G)$ of G is the space of left-invariant vector fields on G under the Lie bracket.

Consider now the commutative diagram

$$\begin{array}{ccc} \text{Lie}(G) & \hookrightarrow & (\mathcal{X}(G), [-, -]) \\ & \searrow \alpha & \downarrow \text{eval}_1 \\ & & T_1(G) \end{array} .$$

Proposition 5.0.3. α is an isomorphism of vector spaces.

Example 5.0.4. Let $G = \text{GL}(n, \mathbb{C})$, which is a complex Lie group. We have that $\text{GL}(n, \mathbb{C})$ is an open submanifold of the vector space $M_n(\mathbb{C})$. Hence $\text{Lie}(G)$ is isomorphic to $M_n(\mathbb{C})$ under the *commutator bracket*, which is given by $[A, B] = AB - BA$.

Definition 5.0.5 (Matrix exponential). Define the map $e^{(\cdot)} : M_n(\mathbb{C}) \rightarrow \text{GL}_n(\mathbb{C})$ by

$$e^A = \sum_{n=0}^{\infty} \frac{A^n}{n!}.$$

This is well-defined. Indeed, letting $\|\cdot\|$ denote the operator norm, we see that $\frac{\|A^n\|}{n!} \leq \frac{\|A\|^n}{n!}$ on any bounded subset $S \subset \mathbb{C}^n$. But $\sum_{n=0}^{\infty} \frac{\|A^n\|^n}{n!} = e^{\|A\|}$ on S , and thus e^A converges uniformly on S . Moreover, one can show that its limit must be invertible.

Exercise 5.0.6. Let $G = \text{SL}_2(\mathbb{C})$, which is complex Lie group. Show that

$$\text{Lie}(G) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(\mathbb{C}) \mid a + d = 0 \right\}.$$

Proof. Any element X of $\text{Lie}(G)$ generates a local flow $\theta : D \subset \mathbb{R} \times G \rightarrow G$. Since X is left-invariant, it is complete. In particular, the maximal integral curve θ^1 is defined on \mathbb{R} . Left-invariance also implies that for any $s \in \mathbb{R}$, $L_{\theta^1(s)} \circ \theta^1$ is an integral curve starting at $\theta^1(s)$. But the curve given by $t \mapsto \theta^1(s+t)$ is also an integral curve starting at $\theta^1(s)$. Hence $\theta^1(s+t) = \theta^1(s)\theta^1(t)$. By the uniqueness of maximal integral curves, this proves that $\theta^1(s)$ is a smooth group homomorphism $\mathbb{R} \rightarrow G$, known as a *one-parameter subgroup* of G .

Moreover, any one-parameter subgroup γ of G has the form $\gamma(t) = e^{tA}$ where $A = \gamma'(0) \in T_1(G) \subset T_1(\mathrm{GL}_2(\mathbb{C})) \cong M_2(\mathbb{C})$. It follows that

$$\begin{aligned} X \in T_1(G) &\iff \forall t \in \mathbb{R}, e^{tX} \in G \\ &\iff \forall t \in \mathbb{R}, \det(e^{tX}) = 1 \\ &\iff \forall t \in \mathbb{R}, e^{t \operatorname{tr}(X)} = 1 \\ &\iff \forall t \in \mathbb{R}, t \operatorname{tr}(X) = 0 \\ &\iff \operatorname{tr}(X) = 0. \end{aligned}$$

□

Remark 5.0.7. Intuitively, Theorem 4.2.15 means that the space $\mathcal{A}^{p,q}(X)$ has a symmetry encoded in the $\mathrm{SL}_2(\mathbb{C})$ -action.

5.1 Lecture 13

Definition 5.1.1. Let V be a vector space endowed with a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$. The *orthogonal group* $\mathrm{O}(V, \langle \cdot, \cdot \rangle)$ is the group of all linear maps $f : V \rightarrow V$ such that $\langle fx, fy \rangle = \langle x, y \rangle$ for any $x, y \in V$.

Example 5.1.2. Consider the Lie group $G := \mathrm{O}(\mathbb{R}^n)$. Define the smooth map $\varphi : \mathrm{GL}_n(\mathbb{R}) \rightarrow \mathrm{GL}_n(\mathbb{R})$ by $A \mapsto AA^t$, which has constant rank. Then $G = \varphi^{-1}(I_n)$, so that $T_{I_n}G = \ker d\varphi_{I_n}$. Since $d\varphi_{I_n}(A) = A^t + A$ for any $A \in M_n(\mathbb{R})$, it follows that $\mathrm{Lie}(G)$ consists of all $n \times n$ skew-symmetric matrices.

Let $(V, \langle -, - \rangle)$ be a finite-dimensional inner product space over \mathbb{R} . For each non-zero $v \in V$, let $\rho_v : V \rightarrow V$ denote the reflection about the hyperplane perpendicular to v , i.e.,

$$\rho_v(w) \equiv w - \frac{2\langle w, v \rangle}{\langle v, v \rangle}v.$$

Note that this is an isometry.

Definition 5.1.3. A finite subset $R \subset V$ with $0 \notin R$ is a *root system* for V if

- (i) $\operatorname{span}(R) = V$,
- (ii) if $a \in R$ and $\lambda r \in R$, then $\lambda = \pm 1$,
- (iii) $\rho_v(a) \in R$ for all $v, a \in R$, and
- (iv) $\frac{2\langle w, v \rangle}{\langle v, v \rangle} \in \mathbb{Z}$ for all $w, v \in R$.

The *rank of R* is $\dim(V)$.

Terminology. An element of R is called a *root*.

An *isomorphism* $R \xrightarrow{\cong} R'$ of root systems is a vector space isomorphism $\varphi : V \xrightarrow{\cong} V'$ such that

$$\begin{aligned} \varphi(R) &= R' \\ \frac{2\langle b, a \rangle}{\langle a, a \rangle} &= \frac{2\langle \varphi(b), \varphi(a) \rangle}{\langle \varphi(a), \varphi(a) \rangle}. \end{aligned}$$

for any $a, b \in R$.

The following result flows easily from the geometric definition of an inner product along with the equality condition of the Cauchy-Schwarz inequality.

Lemma 5.1.4. *Let $a, b \in R$ such that $a \neq \pm b$. Then*

$$\left(\frac{2\langle a, b \rangle}{\langle a, a \rangle} \right) \left(\frac{2\langle b, a \rangle}{\langle b, b \rangle} \right) \in \{0, 1, 2, 3\}.$$

We say that R is *irreducible* if it cannot be partitioned into two nonempty subsets R_1 and R_2 such that $\langle a, b \rangle = 0$ for any $a \in R_1$ and $b \in R_2$.

Proposition 5.1.5. *Every root system can be decomposed into finitely many irreducible root systems.*

Definition 5.1.6. Let R be a root system for V . A subset $B \subset R$ is a *base* for R if

- (i) B is a basis for V and
- (ii) every element of R can be written as an B -combination with coefficients either all positive or all negative.

Proposition 5.1.7. *Every root system has a base.*

Definition 5.1.8 (Dynkin diagram). Let $B := \{b_1, \dots, b_n\}$ be a base for R . The *Dynkin diagram* of R (with respect to B) is the chain graph with vertex set B and exactly

$$\delta_{i,j} := \left(\frac{2\langle b_i, b_j \rangle}{\langle b_i, b_i \rangle} \right) \left(\frac{2\langle b_j, b_i \rangle}{\langle b_j, b_j \rangle} \right) \in \{0, 1, 2, 3\}$$


edges between b_i and b_j ($i \neq j$) such that if $\|b_i\| > \|b_j\|$, then these edges are directed from b_i to b_j .

Terminology. The same graph with only undirected edges is called the *Coxeter graph* of R .

If there is a single edge between two nodes, then it must be undirected. If there are multiple edges between two nodes, then they must be directed.

Theorem 5.1.9.

1. *There is a one-to-one correspondence between (isomorphism classes of) irreducible root systems and finite-dimensional simple Lie algebras over \mathbb{C} .*
2. *There is a one-to-one correspondence between irreducible root systems and Dynkin diagrams of the following forms:*



In particular, we have the following characterizations of classical groups:

$$\begin{aligned} \text{Lie}(\text{SL}_{n+1}(\mathbb{C})) &\longleftrightarrow A_n \\ \text{Lie}(\text{SO}_{2n+1}(\mathbb{C})) &\longleftrightarrow B_n \\ \text{Lie}(\text{Sp}_{2n}(\mathbb{C})) &\longleftrightarrow C_n \\ \text{Lie}(\text{SO}_{2n}(\mathbb{C})) &\longleftrightarrow D_n \quad n \geq 2. \end{aligned}$$

As a result of Theorem 5.1.9, one can show that the following classes of Lie groups are determined by the preceding Dynkin diagrams:

- compact Lie groups,
- simply connected complex Lie groups,
- simply connected compact Lie groups,
- adjoint complex Lie groups, and
- adjoint compact Lie groups.

Example 5.1.10. All of the following Lie groups are determined by A_n .

- $\text{SU}_{n+1}(\mathbb{C}) := \{X \in \text{SL}_{n+1}(\mathbb{C}) \mid XX^* = I_{n+1}\}$
- $\text{SL}_{n+1}(\mathbb{C})$
- $\text{SU}_{n+1}(\mathbb{C})$

$$(d) \text{PSL}_{n+1}(\mathbb{C}) := \frac{\text{SL}_{n+1}(\mathbb{C})}{\mathbb{Z}(\text{GL}_{n+1}(\mathbb{C}) \cap \text{SL}_{n+1}(\mathbb{C}))} \cong \frac{\text{SL}_{n+1}(\mathbb{C})}{\mathbb{Z}/(n+1)}$$

$$(e) \text{PSU}_{n+1}(\mathbb{C})$$

Definition 5.1.11.

1. A *representation* of a Lie group G is a vector space V together with a group homomorphism $G \rightarrow \text{Aut}(V)$.
2. A *representation* of a Lie algebra \mathfrak{g} on a vector space V is a Lie algebra homomorphism $\mathfrak{g} \rightarrow \text{End}(V)$.

A Lie group representation $f : G \rightarrow \text{Aut}(V)$ is *irreducible* if there is no non-trivial subspace of V invariant under $f(g)$ for all $g \in G$. An *irreducible Lie algebra representation* is defined similarly. As it turns out, Theorem 5.1.9 gives rise to a classification of all irreducible representations of complex Lie algebras.

Note 5.1.12. If (V_1, f_1) and (V_2, f_2) are representations of G , then so are both the direct sum $(V_1 \oplus V_2, f_1 \oplus f_2)$ and the tensor product $(V_1 \otimes V_2, f_1 \otimes f_2)$ where

$$\begin{aligned} (f_1 \oplus f_2)(g) &\equiv f_1(g) \oplus f_2(g) \\ (f_1 \otimes f_2)(g) &\equiv f_1(g) \otimes f_2(g) \quad g \in G. \end{aligned}$$

This property also holds for representations of Lie algebras.

For example, suppose that V is a defining representation of dimension two (over \mathbb{C}) with basis $\{x, y\}$. Let $S^k V$ denote a $(k+1)$ -dimensional vector space ($k \geq 0$) with basis $\{x^k, x^{k-1}y, \dots, xy^{k-1}, y^k\}$. Then the tensor product $V^{\otimes k}$ decomposes as $S^k V \oplus \Lambda^k V$.

Now consider the Lie algebra $\text{Lie}(\text{SL}(2))$, which is three-dimensional with basis elements

$$e \equiv \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad f \equiv \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \quad h \equiv \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

such that

$$[e, e] = 0$$

$$[h, e] = 2e$$

$$[f, h] = 2f$$

$$[e, f] = h$$

$$\begin{array}{c|cc} & x & y \\ \hline e & 0 & x \\ f & y & 0 \\ h & x & -y \end{array} \quad x < y$$

Table 1: Action of $\text{Lie}(\text{SL}(2))$ on V

	x^k	$x^{k-1}y$	\dots	y^k
e	0	x^k	\dots	kxy^{k-1}
f	$kx^{k-1}y$	$(k-1)x^{k-2}y^2$	\dots	0
h	kx^k	$(k-2)x^{k-1}y$	\dots	$-ky^k$

Table 2: Action of $\text{Lie}(\text{SL}(2))$ on S^kV .

Observe that e , f , and h act as a lowering, raising, and \pm -grading operators, respectively.

Theorem 5.1.13. *Any finite-dimensional representation of $\text{Lie}(\text{SL}(2))$ is isomorphic to a direct sum of the form*

$$\bigoplus_{k \in \mathbb{Z}_{\geq 0}} S^kV^{\otimes m_k}, \quad m_k \geq 0.$$

For example, we have that

$$\begin{aligned} V \otimes V &\cong S^2V \oplus \Lambda^2V \cong S^2V \oplus S^0V \\ V \otimes S^kV &\cong S^{k+1}V \oplus S^{k-1}V \\ S^kV \otimes S^lV &\cong S^{k+l}V \oplus S^{k+l-2}V \oplus \dots \oplus SV. \end{aligned}$$

Theorem 5.1.14.

1. *If G is a compact Lie group, then any finite-dimensional representation of G is isomorphic to a direct sum of irreducible representations of G .*
2. *If \mathfrak{g} is a semisimple Lie algebra, then any finite-dimensional representation of \mathfrak{g} is isomorphic to a direct sum of irreducible representations of \mathfrak{g} .*

Definition 5.1.15. A *torus* in a compact Lie group G is a connected abelian subgroup of G .

Proposition 5.1.16 (Uniqueness of maximal torus). *Any two maximal tori in G are conjugate in G .*

Proposition 5.1.17. *The rank of a connected compact Lie group G is the dimension of a maximal (w.r.t. inclusion) torus in G .*

Example 5.1.18.

1. The maximal torus in $U(n)$ is precisely the group $U(1)^n \cong (S^1)^n$ of all matrices of the form

$$\begin{bmatrix} S^1 & & 0 \\ & \ddots & \\ 0 & & S^1 \end{bmatrix}.$$

2. The maximal torus in $SU(n)$ has dimension $n - 1$.

In representation theory, we assign to any Lie algebra \mathfrak{g} a lattice $\Lambda_{\mathfrak{g}}$ along with a finite group (or “weight”) W , known as the *Weyl group* of \mathfrak{g} , acting on $\Lambda_{\mathfrak{g}}$. As it turns out, irreducible representations of \mathfrak{g} are in one-to-one correspondence with quotients of the form $\Lambda_{\mathfrak{g}}/W$.

Example 5.1.19. Assign the group $W := \mathbb{Z}/2$ to $\text{Lie}(\text{SL}(2))$. Embed the lattice $\Lambda := \mathbb{Z}$ into $\mathbb{Z} \oplus \mathbb{Z}$ via the mapping $n \mapsto (n, -n)$. This yields an irreducible representation $\Lambda/W \cong \mathbb{Z}_{\geq 0}$ of $\text{Lie}(\text{SL}(2))$.